

THE MERKURIEV-SUSLIN THEOREM FOR ANY SEMI-LOCAL RING

RAYMOND T. HOOBLER

ABSTRACT. We introduce here a method which uses étale neighborhoods to extend results from smooth semi-local rings to arbitrary semi-local rings A by passing to the henselization of a smooth presentation of A . The technique is used to show that étale cohomology of A agrees with Galois cohomology, the Merkuriev-Suslin theorem holds for A , and to describe torsion in $K_2(A)$.

We introduce here a method which uses étale neighborhoods to extend results from smooth semi-local rings to arbitrary semi-local rings. Three applications are given. In the first and last, A is a connected, semi-local ring containing a field k while the second application holds for any connected, semi-local ring A .

1) Let $X = \text{Spec}(A)$, and let n an integer with $(n, \text{char}(k)) = 1$. If F is a finite, locally constant sheaf of $\mathbb{Z}/n$ -modules for the étale site on X , we show

$$H^p(G(A_s/A), F(A_s)) \xrightarrow{\approx} H^p(X, F)$$

where A_s is the separable closure of A , the left hand side is the Galois cohomology of A with coefficients in the $G(A_s/A)$ -module $F(A_s)$ and the right hand side is the étale cohomology group of the semi-local scheme X with coefficients in F .

2) We extend the Merkuriev-Suslin theorem to a connected, semi-local ring A ; that is, for n relatively prime to the residue characteristics of A , the Galois symbol map

$$K_2(A)/n \xrightarrow{\approx} H^2(A, \mathbb{Z}/n(2))$$

is an isomorphism where, as usual, $\mathbb{Z}/n(i) = \mu_n^{\otimes i}$. Since this implies the cup product map is surjective, we conclude that any Azumaya algebra of order n in $Br(A)$ is similar to a tensor product of symbol algebras if A contains a primitive n^{th} root of unity.

3) We extend Suslin's computation of the ℓ -primary component of the torsion in $K_2(k)$, k a field, to A .

Fix notation as follows. $H^p(X, F)$ (or, if $X = \text{Spec}(A)$, $H^p(A, F)$) denotes the étale cohomology group of X with coefficients in the étale

sheaf F . For connected A , Galois cohomology will be indicated by $H^p(\pi_1(A, x), M)$ where $\pi_1(A, x)$ is the algebraic fundamental group of A with base point $x : \text{Spec}(k_s) \rightarrow \text{Spec}(A)$ or, equivalently, by $H^p(G(A_s/A), M)$ ($\pi_1(A, x) = G(A_s/A)$, the Galois group of the separable closure of A , where the base point is defined by embedding the residue field of A into a separably closed field) when M is an abelian group equipped with a continuous action. $X^{(d)}$ stands for the set of generic points x of irreducible components of X of codimension d . If M is a fixed abelian group, we let M/n stand for M/nM , $M\{\ell\}$ stand for the ℓ -primary component of M , and $\text{Div}(M)$ stand for the maximal divisible subgroup of M .

I would like to thank Srinivas, Bruno Kahn, and Wayne Raskind for several very helpful discussions on these and related matters and Chuck Weibel for the last application and unending patience. I also want to express my delight and appreciation to the Tata Institute of Fundamental Research for providing such a fine location and so much stimulation.

1. LOCAL FUNCTORS

Let (A, I) be a pair consisting of a commutative ring with 1 and an ideal $I \subset A$. An étale neighborhood of (A, I) is a pair (A', I') and an étale map $u : A \rightarrow A'$ such that $u(I)A' = I'$ and u induces an isomorphism $\bar{u} : A/I \rightarrow A'/I'$. Geometrically étale neighborhoods of a closed set $W \subset X = \text{Spec}(A)$ look like a cartesian diagram where π is étale:

$$\begin{array}{ccc} W' & \subset & X' \\ \downarrow \approx & & \downarrow \pi \\ W & \subset & X \end{array}$$

The set of all étale neighborhoods of I in A is a filtered category which we denote $\text{Et}(I)$. If we assume that $I \subseteq \text{rad}(A)$, then $\text{Et}(I)$ is used to define the *henselization* of the pair (A, I) [15] by setting

$$A_{/I}^h := \varinjlim_{(A', I') \in \text{Et}(I)} A'.$$

Then the pair $(A_{/I}^h, I^h)$ is a hensel pair where $I^h = IA_{/I}^h$. If $I = \mathfrak{m}$ is maximal, we get the hensel local ring $A_{\mathfrak{m}}^h$.

Let $\mathcal{C}$ be a category containing $\text{Et}(I)$ as a full subcategory and $A_{/I}^h$. We introduce the definition of a *local* functor.

Definition 1. A covariant functor $F : \mathcal{C} \rightarrow \text{Ab}$ is said to be *local* (for the étale topology) with respect to a closed set $V(I)$ in $\text{Spec}(A)$ if

(1) I is a radical ideal in A , $Et(I)$ is a full subcategory of $\mathcal{C}$, and $A^h/I \in \mathcal{C}$

(2) the natural map

$$(1) \quad \varinjlim_{(A', I') \in Et(I)} F(A') \rightarrow F(A/I)$$

is an isomorphism.

We say that F is local for the étale topology on X if, for all affine open subsets $Spec(A) \subset X$ and all closed sets $V(I) \subset Spec(A)$, F is local with respect to the closed set $V(I) \subset Spec(A)$. In many cases of importance the limit condition above can be replaced with a condition involving the henselization of A along I . Recall that a covariant functor $F : \mathcal{C} \rightarrow Ab$ is said to be locally of finite presentation if for any filtered inductive limit $A = \varinjlim_{i \in I} A_i$, the natural map

$$\varinjlim_{i \in I} F(A_i) \rightarrow F(A)$$

is an isomorphism.

For such functors we are only interested in a *hensel pair condition*.

Definition 2. A functor $F : \mathcal{C} \rightarrow Ab$ is said to satisfy the *hensel pair condition for I* , resp. *epic hensel pair condition for I* , if $F(A) \rightarrow F(A/I)$ is an isomorphism, resp. an epimorphism, for any hensel pair (A, I) . The functor satisfies the *hensel pair condition*, resp. *epic hensel pair condition* if it does so for any hensel pair $(A, I) \in \mathcal{C}$.

Then if F is locally of finite presentation and F satisfies the hensel pair condition, F is local for the closed set $V(I)$. Thus in applying this definition we will first verify that F is locally of finite presentation and then that $F(A^h) \rightarrow F(A^h/I^h)$ is an isomorphism when (A^h, I^h) is a hensel pair.

We are primarily interested in three examples of functors satisfying the hensel pair condition.

Example 1.

- (1) $H^i(-, F)$ where F is a locally constant sheaf of $\mathbb{Z}/n$ modules on $Spec(A)_{et}$ ([18] or [5])
- (2) $K_i(-; \mathbb{Z}/n)$ where n is relatively prime to the residue characteristics of A ([4]) and
- (3) $K_2(-)/n$ where n is relatively prime to the residue characteristics of A .

The last example is easily seen to satisfy the hensel pair condition. Surjectivity of $K_2(A)/n \rightarrow K_2(A/I)/n$ for a hensel pair (A, I) follows

immediately since K_2 is generated by symbols, and injectivity follows by a straightforward calculation done in [11, Appendix] or from Gabber's theorem [4]. Alternatively we could use the universal coefficient theorem, the second example, and note that the third term in the universal coefficient theorem is the n -torsion in K_1 which is the n^{th} roots of unity. These examples will be discussed at greater length in the applications.

Our applications are a result of the following theorem.

Theorem 1. *Let A be a semi-local ring essentially of finite type over a base ring k . Suppose $A \cong B/J$ where B is a smooth semi-local ring essentially of finite type over k . Let C be a category of semi-local ring extensions of B containing A , $B_{/J}^h$, and the full subcategory $Et(J)$. Suppose $F_1, F_2 : C \rightarrow Ab$ are two covariant functors which are either both local for $V(J)$ or F_2 is local for $V(J)$ and F_1 is locally of finite presentation and satisfies the epic hensel pair condition. Let $\phi : F_1 \rightarrow F_2$ be a natural transformation such that $\phi(B')$ is an isomorphism if B' is smooth and essentially of finite type over k . Then $\phi(A)$ is an isomorphism.*

Proof. If A is essentially of finite type over k , then A has a presentation as B/J where B is essentially of finite type and smooth over k . Thus $\phi(B)$ is an isomorphism as is $\phi(B')$ for any $B' \in Et(J)$ and so, by (1), $\phi(A)$ must be an isomorphism in the first case. In the second case, $F_1(B_{/J}^h) \rightarrow F_1(A)$ must be an isomorphism since F_2 is local for $V(J)$ and $\phi(B_{/J}^h)$ is an isomorphism. $\square$

Corollary 1. *In the above situation F_1 is local for $V(J)$ if it is locally of finite presentation and satisfies the epic hensel pair condition.*

2. APPLICATIONS

We have three applications of this perspective. They all rely on first establishing an isomorphism for semi-local rings smooth over a base scheme and then using the examples above to extend the result to arbitrary semi-local schemes over the base.

2.1. Galois cohomology = Étale cohomology. As a first application we consider the relationship between étale and Galois cohomology of semi-local rings. While it is a generally accepted fact for smooth local rings over a field, the details do not, as far as I know, appear in the literature although there is an argument, due to Bruno Kahn, when the generalized Kato conjecture holds [11]. We begin by reviewing and

reinterpreting Galois cohomology and then proving the isomorphism in this case.

Let S be an arbitrary connected scheme. Define a site S_{ret} by considering the category of schemes $T \rightarrow S$ which are finite and étale (= revêtement étale) over S . A covering morphism is a surjection of schemes over S , $T_2 \rightarrow T_1$, which will necessarily be finite and étale. This is the same as considering the class $E = (ret)$ of all finite, étale morphisms [12, Chapter II, Section 1]. S_{ret} is then the small site $(E/S)_E$ where E consists of "revêtements étales" as above. This site is discussed in [12, Chapter 1] where it is called FEt and used to describe the fundamental group of S . Cohomology in S_{ret} can be identified with Galois cohomology according to the following recipe.

Suppose F is a sheaf on S_{ret} . Then F takes finite, disjoint unions of schemes to direct products. Thus the discussion in [12, Chapter III, Example 2.6] applies to the covering $T \rightarrow S$ in S_{ret} where T is Galois over S with group G . Hence

$$(2) \quad \check{H}^p(T/S, F) \cong H^p(G, F(T))$$

where $F(T)$ is a left G -module via the action of G on T .

Let $\iota : \mathbf{S}(S_{ret}) \rightarrow \mathbf{P}(S_{ret})$ be the forgetful functor that regards a sheaf as a contravariant functor defined on S_{ret} . When we pass to the limit over all coverings T/S in S_{ret} , the left hand side of (2) becomes $\check{H}^p(S_{ret}, \iota F)$. If we fix a geometric point $x : Spec(k_s) \rightarrow S$ where k_s is a separably closed field, then the theory of the fundamental group expresses any finite, étale covering $T \rightarrow S$ as a quotient of a Galois covering. Consequently the right hand side of (2) is $\varinjlim H^p(G_i, F(T_i))$ where G_i ranges through the finite quotients of $\pi_1(S, x)$ and $T_i \rightarrow S$ is the corresponding étale covering with group G_i . Let $F(S_s)$ denote the abelian group $\varinjlim F(T_i)$. If F is a locally constant sheaf on S , then the subgroup of $\pi_1(S, x)$ defining the covering $T \rightarrow S$ such that $F|_T$ is constant is of finite index in $\pi_1(S, x)$ and acts trivially on $F(S_s)$. Hence $F(S_s)$ has a continuous $\pi_1(S, x)$ -action. In general this need not be the case, but we have enough to conclude that for F a locally constant sheaf on S_{ret} ,

$$(3) \quad H^p(S_{ret}, F) \cong H^p(\pi_1(S, x), F(S_s))$$

(Note that a finite, locally constant sheaf for the étale topology is the same as a finite, locally constant sheaf on S_{ret} .) In view of the isomorphism (2) on S_{ret} , this is essentially the statement that sheaf cohomology coincides with Čech cohomology. Corollary 2.5 of Chapter III [12] gives sufficient conditions for this; namely, for every surjection

$F \rightarrow F''$ of sheaves, the map

$$\varinjlim \prod F(U_{i_0 \dots i_p}) \rightarrow \varinjlim \prod F''(U_{i_0 \dots i_p})$$

is surjective where the limit is over all coverings $\{\coprod U_i \rightarrow S\}$ of S and $U_{i_0 \dots i_p} = U_{i_0} \times_S \dots \times_S U_{i_p}$. But if $T \rightarrow S$ is a Galois covering of S with group G , then $F(T_{i_0 \dots i_p}) = F(\coprod_{G \times p} T) = \prod_{G \times p} F(T)$. Thus taking limits of $F(T_{i_0 \dots i_p})$ over all coverings means taking limits over all coverings of products of copies of $F(T)$ and so surjectivity is immediate.

This discussion reduces our first application to the following theorem about the change of sites morphism $\tau : S_{et} \rightarrow S_{ret}$.

Theorem 2. *Let A be a connected semi-local ring containing a field k . If F is a finite, locally constant sheaf of $\mathbb{Z}/n$ -modules for the étale site on A where $(n, \text{char}(k)) = 1$, then*

$$(4) \quad H^p(A_{ret}, \tau_* F) \rightarrow H^p(A_{et}, F)$$

is an isomorphism for all p .

Proof. We wish to apply our extension theorem. We begin by treating the case of a smooth, semi-local ring R essentially of finite type over k , an algebraically closed field first. We will show that $R^q \tau_* F = 0$ for $q > 0$ and any finite, locally constant sheaf F of $\mathbb{Z}/n$ -modules. Suppose $x \in H^q(R, F)$ with $q > 0$. We may assume $F = C_X$ is a constant sheaf of $\mathbb{Z}/n$ -modules with value C since F becomes constant after a finite, étale extension of R . By assumption there is a smooth, connected variety X over k , closed points $t_i \in X$, $1 \leq i \leq m$, such that $R \cong O_{X, t_1 \dots t_m}$ and an element $x' \in H^q(X, C_X)$ such that $x' |_{\text{Spec}(O_{X, t_1 \dots t_m})} = x$. We construct a finite, étale covering $\rho : X' \rightarrow X$ such that $\rho^*(x') = 0$ using induction on $\dim(X)$ and the existence of Artin neighborhoods. If $\dim(X) = 0$, the assertion is obvious. If $\dim(X) = d$, the existence of an Artin neighborhood relative to k means there is a diagram ([12, p. 117] or [16])

$$\begin{array}{ccccc} X & \xrightarrow{j} & \overline{X} & \xleftarrow{i} & Y \\ & f \searrow & \downarrow \overline{f} & \swarrow g & \\ & & X_0 & & \end{array}$$

in which

- (1) j is an open immersion, dense in each fibre and $Y = \overline{X} - X$;
- (2) $\overline{f}$ is smooth and projective with geometrically irreducible fibres of dimension one;
- (3) g is finite and étale and each fibre of g is non-empty.

(Although the statement of the existence of Artin neighborhoods given in [12] refers only to a local ring, the proof as given in [16, Exposé

XI, Section 3] clearly extends to the semi-local case. In fact, the proof of Proposition 3.3, the key statement, is footnoted to that effect. Since base extension of a local ring from an arbitrary field k to its separable closure k_s may produce a semi-local ring we need this more general result.)

Now $(Y, \overline{X})$ is a smooth X_0 -pair. The Gysin sequence [12, Chapter VI, Corollary 5.3 and Remark 5.4 (a)] becomes $\overline{f}_* C_{\overline{X}} \cong f_* C_X$,

$$0 \rightarrow R^1 \overline{f}_* C_{\overline{X}} \rightarrow R^1 f_* C_X \rightarrow g_*(C_Y \otimes T_{Y/\overline{X}}) \rightarrow R^2 \overline{f}_* C_{\overline{X}} \rightarrow R^2 f_* C_X \rightarrow 0,$$

and $R^p \overline{f}_* C_{\overline{X}} \cong R^p f_* C_X$ for $p > 2$. The base change maps for the proper morphisms $\overline{f}$ and g are isomorphisms. Consequently the base change map for f is also an isomorphism. Since the fibres of f are non-complete curves, we find $R^p f_* C = 0$ for $p > 1$. Moreover $\overline{f}$ is smooth and so $R^1 \overline{f}_* C_{\overline{X}}$ is finite and locally constant on X_0 as is $g_*(C_Y \otimes T_{Y/X})$ since g is an étale covering space and $T_{Y/X}$ is locally constant on Y . Therefore we conclude that $R^p f_* C_X$ is a finite, locally constant sheaf of $\mathbb{Z}/n$ -modules for $p = 0$ or 1 .

This information shows that the Leray spectral sequence for f degenerates into the long exact sequence

$$(5) \quad \begin{aligned} \dots \rightarrow H^q(X_0, f_* C_X) \rightarrow H^q(X, C_X) \rightarrow H^{q-1}(X_0, R^1 f_* C_X) \\ \rightarrow H^{q+1}(X_0, f_* C_X) \rightarrow \dots \end{aligned}$$

and we may apply the induction hypothesis. Then for any

$$y \in H^{q-1}(X_0, R^1 f_* C_X),$$

there is, after shrinking X_0 if necessary, a finite, étale covering space $\rho' : X'_0 \rightarrow X_0$ such that $\rho'^*(y) = 0$. Now by using the pullbacks of appropriate coverings of X_0 to X and a diagram chase, it is a straightforward matter to produce a finite, étale covering $\rho : X' \rightarrow X$ such that $\rho^*(x') = 0$.

This argument is, of course, the essence of Remark 3.16 [12, Chapter III] and shows that $R^q \tau_* F = 0$ for $q > 0$. Hence the Leray spectral sequence for the change of site morphism $\tau : X_{et} \rightarrow X_{ret}$ collapses, and we conclude

$$H^p(R_{ret}, \tau_* F) \cong H^p(R_{ét}, F)$$

Now suppose that R is a semi-local ring which is smooth and essentially of finite type over k , a not necessarily algebraically closed field. Then for any scheme X there is a Hochschild-Serre spectral sequence [12, Chapter III, Remark 2.21(b)]

$$(6) \quad H^p(G(k_s/k), H^q(X_s, F)) \Rightarrow H^n(X, F)$$

where $X_s = X \times_k \text{Spec}(k_s)$. Similarly the theory of algebraic fundamental groups constructs a profinite group extension

$$1 \rightarrow G((R \otimes_k k_s)_s / R \otimes_k k_s) \rightarrow G(R_s / R) \rightarrow G(k_s / k) \rightarrow 1$$

and so there is a corresponding Hochschild-Serre spectral sequence

$$(7) \quad \begin{aligned} H^p(G(k_s/k), H^q(G((R \otimes_k k_s)_s / R \otimes_k k_s), F((R \otimes_k k_s)_s))) \Rightarrow \\ H^n(G(R_s/R), F(R_s)) \end{aligned}$$

The change of site morphism defines a homomorphism Ψ from (6) to (7).

Since F is a sheaf of $\mathbb{Z}/n$ -modules where n is relatively prime to $\text{char}(k)$, we conclude, as usual, that $H^p(X \times_k \text{Spec}(k_s), F) \cong H^p(X \times_k \text{Spec}(\bar{k}), F)$ where $\bar{k}$ is the algebraic closure of k . Consequently, by the first case, Ψ is an isomorphism on the $E_2^{p,q}$ terms. This shows that (4) is an isomorphism when A is essentially smooth over a not necessarily algebraically closed field k .

Some preparation is required in order to apply Theorem 1 to an arbitrary semi-local ring A whether we work with the small or large étale site. Our hypothesis on F shows that F is representable in either the small or large site by a semi-local scheme $\mathbb{F}$ which is finite and étale over A . Now A is a filtered limit of connected semi-local rings A_i essentially of finite type over k . Consequently there is a finite, étale covering space $\mathbb{F}_{i_0}$ over A_{i_0} for a sufficiently large A_{i_0} such that $\mathbb{F}_{i_0} \times_{A_{i_0}} A \cong \mathbb{F}$. We use $\mathbb{F}_{i_0}$ to define a locally constant sheaf F_{i_0} over A_{i_0} whose restriction to A is F . Next find a presentation of A_{i_0} by a smooth connected semi-local ring B_{i_0} essentially of finite type over k so that $A_{i_0} \cong B_{i_0}/I$. Since any étale covering space of A_{i_0} may be lifted to an étale covering space of B_{i_0} (by the theorem of the primitive element all one has to do is lift a separable polynomial), we may assume there is a finite, locally constant sheaf G on $\text{Spec}(B_{i_0})$ whose restriction to $\text{Spec}(A)$ is F .

Let C be the category of connected semi-local rings over B , $F_1 = H^p(-_{ret}, \tau_* G)$, and $F_2 = H^p(-, G)$ where now we consider G on the large site over $\text{Spec}(B)$. Note that G is represented by a finite étale covering space B' of B in B_{et} and so G is also represented by B' on the big étale site over B . Since B' also defines a covering in B_{ret} the same observation applies to the sheaf $\tau_* G$ in the big site B_{ret} . After showing that the hypotheses of our extension theorem are met we can conclude that (4) is an isomorphism for any B -algebra A with the property that $G|_A = F$ and $\tau_* G|_A = \tau_* F$.

Both F_1 and F_2 are locally of finite presentation. For F_2 this is [12, Chapter III, Lemma 1.16] suitably interpreted. For F_1 we need

a different argument. Suppose $R = \varinjlim R_i$ with R_i a connected semi-local ring over B where $i \in I$, a filtered set. Any étale covering space of $\varinjlim R_i$ comes from an étale covering space defined over one of the R_i and homomorphisms between any two étale covering spaces of $\varinjlim R_i$ descend to a homomorphism between covering spaces over one of the R_i . Thus we have an isomorphism

$$\lim_{R_i} \lim_{T_i/R_i} \check{H}^p(T_i/R_i, G) \rightarrow \lim_{T/R} \check{H}^p(T/R, G)$$

since the Čech cohomology groups depend only on the Galois group of the covering by (2) and G is locally constant. Thus, using the Galois cohomology interpretation of F_1 , we also have

$$H^p(R_{ret}, \tau_* G) \cong \varinjlim H^p(R_{i ret}, \tau_* G)$$

and so F_1 is locally of finite presentation.

Now we turn to the hensel pair condition. Suppose (R, J) is a hensel couple and G is a locally constant sheaf of $\mathbb{Z}/n$ -modules on $\text{Spec}(R)$. Then G is represented by a finite, étale group scheme over R , and, by Example 1, we have an isomorphism for a hensel couple (R, J) and such a G ,

$$H^p(R_{ét}, G) \cong H^p(R/J_{ét}, G|_{R/J}).$$

Thus F_2 is a local functor for the étale topology satisfying the hensel pair condition. As for F_1 , there is a one-to-one correspondence between étale covering spaces of R/J and étale covering spaces of R , and so we have an isomorphism

$$H^p(R_{ret}, \tau_* G) \cong H^p(R/J_{ret}, \tau_*(G|_{R/J})).$$

Hence F_1 is a local functor for the étale topology and also satisfies the hensel pair condition. We can now apply Theorem 1 to conclude the proof. $\square$

Corollary 2. *Let A be a semi-local ring containing a field k . Suppose F is a finite, locally constant sheaf of $\mathbb{Z}/n$ -modules where $(n, \text{char}(k)) = 1$. Then for any $x \in H^p(A, F)$, $p > 0$, there is a Galois extension A'/A such that $x|_{A'} = 0 \in H^p(A', F)$.*

2.2. Merkurjev-Suslin theorem. Our second application is an extension of the Merkurjev-Suslin theorem to an arbitrary semi-local ring.

For a semi-local ring A and any n which is relatively prime to the residue characteristics of A , Kummer theory provides a natural isomorphism $K_1(A)/n \rightarrow H^1(A, \mathbb{Z}/n(1))$. The Galois symbol map [20] is obtained by extending this map multiplicatively to $K_i(A)/n \rightarrow H^i(A, \mathbb{Z}/n(i))$.

Theorem 3. *Let A be a semi-local ring. Then if n is relatively prime to the residue characteristics of A , the Galois symbol map*

$$(8) \quad K_2(A)/n \rightarrow H^2(A, \mathbb{Z}/n(2))$$

is an isomorphism.

Corollary 3. *Let A be a semi-local ring, and suppose n is an integer which is relatively prime to the residue characteristics of A . Then the cup product map*

$$H^1(A, \mathbb{Z}/n(1)) \times H^1(A, \mathbb{Z}/n(1)) \rightarrow H^2(A, \mathbb{Z}/n(2))$$

is onto. In particular, if A contains a primitive n^{th} root of unity, the n -torsion in the Brauer group of A is generated by symbol algebras $(a, b)_n$.

Corollary 4. *Let X be a scheme, and suppose n is an integer which is relatively prime to the residue characteristics of X . The symbol map defines an isomorphism of Zariski sheaves*

$$\underline{K_2/n} \rightarrow \underline{H^2(\mathbb{Z}/n(2))}.$$

The proof requires some preliminary material on K-theory with coefficients, Chern classes, and a discussion of results of Gillet which will be used to prove the theorem for semi-local rings smooth and essentially of finite type over a semi-local Dedekind ring.

Fix a ring A , a prime ℓ that is a unit in A , and an integer n which is at least 2 if $\ell = 2$ (See [21, Proposition 2.4] for a detailed discussion of the case $\ell = 2$.) Algebraic K-theory with coefficients, $K_r(A; \mathbb{Z}/\ell^n)$, is a bifunctor in A and the cyclic group $\mathbb{Z}/\ell^n$, and, for a pair of integers $1 \leq i, 0 \leq j \leq 2i$, there is a theory of Chern classes given by natural transformations

$$c_{i,j}^{\ell^n} : K_{2i-j}(A; \mathbb{Z}/\ell^n) \rightarrow H^j(A, \mathbb{Z}/\ell^n(i))$$

satisfying certain properties. Among the properties these objects satisfy are:

(1) (universal coefficient theorem) There is a natural short exact sequence

$$(9) \quad 0 \rightarrow K_r(A)/\ell^n K_r(A) \rightarrow K_r(A; \mathbb{Z}/\ell^n) \rightarrow {}_{\ell^n}K_{r-1}(A) \rightarrow 0$$

(2) (functoriality) For any $n > m$, the diagram

$$\begin{array}{ccc} K_{2i-j}(A; \mathbb{Z}/\ell^m) & \rightarrow & K_{2i-j}(A; \mathbb{Z}/\ell^n) \\ \downarrow c_{ij}^{\ell^m} & & \downarrow c_{ij}^{\ell^n} \\ H^j(A, \mathbb{Z}/\ell^m(i)) & \rightarrow & H^j(A, \mathbb{Z}/\ell^n(i)) \end{array}$$

defined from the inclusion $\mathbb{Z}/\ell^m \hookrightarrow \mathbb{Z}/\ell^n$ commutes.

- (3) (naturality) For any valuation ring $\mathcal{O}_v$ and prime ℓ distinct from the residue characteristic of $\mathcal{O}_v$ and any n , we have a commutative diagram

$$\begin{array}{ccc} K_{2i-j}(F; \mathbb{Z}/\ell^n) & \xrightarrow{\bar{\partial}} & K_{2i-j-1}(\kappa(v); \mathbb{Z}/\ell^n) \\ \downarrow c_{ij}^{\ell^n} & & \downarrow (1-i)c_{i-1j-1}^{\ell^n} \\ H^j(F, \mathbb{Z}/\ell^n(i)) & \xrightarrow{\partial} & H^{j-1}(\kappa(v), \mathbb{Z}/\ell^n(i-1)) \end{array}$$

where $\kappa(v)$ is the residue field of $\mathcal{O}_v$ and F is its field of fractions. In particular the Chern classes fit together to define a map from the Quillen-Gersten complex for $K_*(-; \mathbb{Z}/\ell^n)$ to the Bloch-Ogus complex for $H^*(-, \mathbb{Z}/\ell^n(\cdot))$. (Quillen's argument in [14] applies unchanged to $K_i(-; \mathbb{Z}/m)$ where $m = \ell^n$.)

Details of the above properties can be found in [17].

Now the other tool we need is a reformulation and extension to a semi-local ring of a result of Gillet-Levine[7] and Gillet[6].

Theorem 4 (Gillet-Levine, Gillet). *Let B be a connected, semi-local ring with quotient field K which is smooth and essentially of finite type over a semi-local Dedekind ring D . Then, for any integer n relatively prime to the residue characteristics of B and any $q \geq 0$, we have a Gersten-Quillen resolution of $K_q(B; \mathbb{Z}/n)$:*

$$0 \rightarrow K_q(B; \mathbb{Z}/n) \rightarrow K_q(K; \mathbb{Z}/n) \rightarrow \coprod_{x \in B^{(1)}} K_{q-1}(\kappa(x); \mathbb{Z}/n) \rightarrow \dots$$

Proof. The argument of Gillet and Levine and Gillet immediately extends to semi-local rings as is clear from going through their arguments. $\square$

We also need a weak version of this result for etale cohomology. This has been demonstrated by Gillet but remains unpublished[8]. A proof for n an odd integer relatively prime to the residue characteristics of B is given in an appendix at the end of this paper.

Lemma 1. *Let B be a connected, semi-local ring which is smooth and essentially of finite type over a semi-local Dedekind ring, $\mathcal{D}$. Let K be the quotient field of B . Then, for any integer n which is relatively prime to the residue characteristic of $\mathcal{D}$,*

$$H^2(B, \mathbb{Z}/n(2)) \rightarrow H^2(K, \mathbb{Z}/n(2))$$

is a monomorphism.

We will need the universal coefficient theorem to produce a Gersten-Quillen sequence for $K_2(-)/n$ using work of Gillet and Gillet-Levine

and then the compatibility of the Chern class map with the Quillen-Gersten complex and the Bloch-Ogus complex will do the rest in the smooth case.

Proof. We deal with a semi-local algebra A over a mixed characteristic Dedekind ring $\mathcal{D}$ first. We may assume A is connected. Both $K_2(-)/n$ and $H^2(-, \mathbb{Z}/n(2))$ are locally of finite presentation over $\text{Spec}(\mathcal{D})$, and so we may assume A is essentially of finite type over $\mathcal{D}$. Now realize $A \cong B/J$ where B is a smooth, semi-local ring which is essentially of finite type over $\mathcal{O}_v$. Let $\mathcal{C}$ be the category of connected semi-local rings essentially of finite presentation over $\mathcal{D}$. We wish to apply Theorem 1 to this situation with $k = \mathcal{D}$, $F_1 = K_2(-)/n$, $F_2 = H^2(-, \mathbb{Z}/n(2))$, and $\phi : F_1 \rightarrow F_2$ being the Galois symbol map.

First observe that $K_2(-)/n$ satisfies the epic hensel pair condition since it is generated by symbols and $H^2(-, \mathbb{Z}/n(2))$ satisfies the hensel pair condition by Gabber or Strano's result ([5] or [18]). It remains to show that $\phi(B)$ is an isomorphism if B is smooth and essentially of finite type over $\mathcal{D}$. The exactness of

$$(10) \quad K_2(B; \mathbb{Z}/n) \hookrightarrow K_2(K; \mathbb{Z}/n) \rightarrow \coprod_{x \in (\text{Spec } B)^{(1)}} K_1(\kappa(x); \mathbb{Z}/n)$$

where K is the quotient field of B was shown by Gillet [6] using work of Gillet-Levine [7] (see Theorem 2.6). We see that $K_1(\kappa(x); \mathbb{Z}/n) \cong K_1(\kappa(x))/n$, $K_2(B; \mathbb{Z}/n)$ is an extension of $\mu_n(B) (= {}_n K_1(B))$ by $K_2(B)/n$, and similarly for $K_2(K; \mathbb{Z}/n)$. We can now reinterpret (10) as the exact sequence of the first line below

$$\begin{array}{ccccc} K_2(B)/n & \hookrightarrow & K_2(K)/n & \rightarrow & \coprod_{x \in (\text{Spec } B)^{(1)}} K_1(\kappa(x))/n \\ \downarrow & & \downarrow \cong & & \downarrow \cong \\ H^2(B, \mathbb{Z}/n(2)) & \hookrightarrow & H^2(K, \mathbb{Z}/n(2)) & \rightarrow & \coprod_{x \in (\text{Spec } B)^{(1)}} H^1(\kappa(x), \mathbb{Z}/n(1)) \end{array}.$$

Here the bottom row is not necessarily exact but is a complex, and the first map is a monomorphism by Lemma 1, the first vertical isomorphism is the Merkuriev-Suslin theorem for fields, and the second vertical isomorphism is the observation that both groups are isomorphic to $\coprod_{x \in (\text{Spec } B)^{(1)}} \kappa(x)^*/\kappa(x)^{*n}$. The diagram commutes by the naturality condition above. Since the bottom row is a complex we conclude that the first vertical map is an isomorphism as desired. Theorem 1.1 now finishes this case.

The case of a semi-local ring containing a field is similar but simpler since we can use Grayson's version of the Gersten-Quillen sequence [9] and the Bloch-Ogus sequence [3] in place of the argument involving algebraic K-theory with coefficients and Lemma 1. $\square$

Suppose the Bloch-Kato conjecture holds for fields; that is, the Galois symbol map

$$(11) \quad K_i^M(K)/n \rightarrow H^i(K, \mathbb{Z}/n(i))$$

is an isomorphism where $K_i^M(K)$ is the i^{th} Milnor K-group of the field K and n is relatively prime to $\text{char}(K)$. Then the same argument applies since the Gersten-Quillen and Bloch-Ogus sequences hold for i as well as 2, at least up to $(i-1)!$ torsion and Gillet's unpublished result. Thus the Bloch-Kato conjecture for fields in degree i and $i-1$ would show that (11) is an isomorphism for smooth local rings essentially of finite type over k . The rest of the argument is then identical, and so we would conclude that $K_i^M(A)/n \rightarrow H^i(A, \mathbb{Z}/n(i))$ is an isomorphism if A is a semi-local ring containing k and $(n, (i-1)!\text{char}(k)) = 1$.

2.3. Torsion in K_2 . The final application was suggested by C. Weibel. Recall that Suslin in [19] used Chern classes to construct an isomorphism

$$K_2(F)\{\ell\} \xrightarrow{\approx} H^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))/\text{Div}(H^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2)))$$

where F is either a field of positive characteristic and ℓ is a prime distinct from $\text{char}(F)$, or F is a field of finite type over $\mathbb{Q}$. Note that Suslin is using continuous Galois cohomology since he uses the identification

$$H^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))/\text{Div}(H^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))) = H_{\text{cont}}^2(F, \mathbb{Z}_\ell(2))\{\ell\}$$

where continuous cochains must be used to correctly identify $H_{\text{cont}}^2(F, \mathbb{Z}_\ell(2))\{\ell\}$ as $H^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$ modulo its maximal divisible subgroup.

Theorem 5. *Let A be a semi-local ring containing a field k and suppose that A is either essentially of finite type over $\mathbb{Q}$ or $\text{char}(k) > 0$. Let ℓ be a prime distinct from $\text{char}(k)$. Then the Chern class $c_{2,1}$ induces an isomorphism*

$$K_2(A)\{\ell\} \xrightarrow{\approx} H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))/\text{Div}(H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))) \xrightarrow{\approx} H_{\text{cont}}^2(A, \mathbb{Z}_\ell(2))\{\ell\}$$

where $H_{\text{cont}}^2(A, \mathbb{Z}_\ell(2))$ is continuous étale cohomology [10].

Corollary 5. *Suppose A is a unibranch, e.g. normal, local ring containing a field k with $\text{char}(k) = p > 0$, and let ℓ be as above. Then*

$$K_2(A)\{\ell\} \xrightarrow{\approx} H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$$

The conclusion of Theorem 5 is independent of the characteristic of k , but the argument required in the two cases is different. We begin by recalling the structure of Suslin's argument in [19] and the role of

continuous cohomology. He first observes (in the proof of Proposition 3.8) that $c_{2,1}$ appears in a commutative, exact diagram

$$(12) \quad \begin{array}{ccccccc} K_3(F) & \rightarrow & K_3(F; \mathbb{Z}/\ell^n) & \xrightarrow{c_{2,1}} & H^1(F, \mathbb{Z}/\ell^n(2)) \\ \downarrow & & & & \downarrow \\ H_{cont}^1(F, \mathbb{Z}_\ell(2)) & \rightarrow & H_{cont}^1(F, \mathbb{Q}_\ell(2)) & \rightarrow & H_{cont}^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2)) & \rightarrow & H_{cont}^2(F, \mathbb{Z}_\ell(2))\{\ell\} \rightarrow 0 \end{array}$$

where the right hand vertical arrow comes from the coefficient sequence defined by multiplication by ℓ^n on $\mathbb{Q}_\ell/\mathbb{Z}_\ell(2)$. Continuous cohomology is used here to make the bottom sequence exact (the first arrow is defined on ℓ -adic sheaves by sending $x \mapsto \ell^{-n}x$) and to identify the image of $H_{cont}^1(F, \mathbb{Q}_\ell(2))$ with the maximal divisible subgroup of $H_{cont}^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$ and the image of $H_{cont}^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$ with the ℓ -primary torsion in $H_{cont}^2(F, \mathbb{Z}_\ell(2))$. Now by exactness and the universal coefficient theorem $c_{2,1}$ factors through $K_2(F)\{\ell\}$ as

$$\begin{aligned} \bar{c}_{2,1} : K_2(F)\{\ell\} &\rightarrow H_{cont}^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))/\text{Image}(H_{cont}^1(F, \mathbb{Q}_\ell(2))) \\ &\cong H_{cont}^2(F, \mathbb{Z}_\ell(2))\{\ell\}. \end{aligned}$$

Suslin has already shown that $H_{cont}^1(F, \mathbb{Z}_\ell(2))$ is finite when $\text{char}(F) > 0$ [19, Corollary 2.8], and so $H_{cont}^1(F, \mathbb{Q}_\ell(2)) = 0$ and

$$H_{cont}^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2)) \cong H_{cont}^2(F, \mathbb{Z}_\ell(2))\{\ell\}$$

in this case. He then shows that $\bar{c}_{2,1} : K_2(F)\{\ell\} \rightarrow H_{cont}^1(F, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$ is an isomorphism [19, Theorem 3.9] by an induction argument. In characteristic zero $H_{cont}^1(F, \mathbb{Q}_\ell(2))$ is only zero if F_0 , the algebraic closure of $\mathbb{Q}$ in F , has only real embeddings in $\mathbb{C}$, but he can reduce the theorem to F_0 [19, Proposition 3.3] and here the desired result was proven earlier by Tate. Thus extending this result to a semi-local ring containing a field which is essentially of finite type over $\mathbb{Q}$ in characteristic zero requires first defining the factorization of $c_{2,1}$ through $\bar{c}_{2,1}$ (which will require continuous étale cohomology [10]) and then showing that $\bar{c}_{2,1}$ is an isomorphism.

$c_{2,1}$ is defined for arbitrary semi-local rings A (at least if $n > 2$ when $\ell = 2$) and Suslin's factorization argument applies which allows us to define

$$(13) \quad \bar{c}_{2,1}(A) : K_2(A)\{\ell\} \rightarrow H_{cont}^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))/\text{Div}^1(A)$$

where, for simplicity, $\text{Div}^1(A)$ denotes the maximal ℓ -divisible subgroup of $H_{cont}^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$. The exact sequence needed for this factorization requires using continuous étale cohomology which Jannsen

[10] has developed. It agrees with continuous group cochain cohomology when A is a field and has all the needed properties to construct the exact sequence in (12) and so to show that $\bar{c}_{2,1}$ is properly defined. Moreover, for a coefficient system of sheaves F_n , when $H^{i-1}(A, F_n)$ satisfies the Mittag-Leffler condition, $H_{cont}^i(A, \varprojlim F_n)$ is the usual ℓ -adic cohomology. Thus $H_{cont}^1(A, \mathbb{Z}_\ell(2)) = H^1(A, \mathbb{Z}_\ell(2))$ and $H_{cont}^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2)) = H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$. In addition, [10, Theorem 5.14] is the identification $H_{cont}^2(A, \mathbb{Z}_\ell(2))\{\ell\} \cong \text{Im}(H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2)))$.

The proof of Theorem 3, while different in $\text{char}(A) > 0$ and $\text{char}(A) = 0$, starts from the comparison of the Gersten-Quillen exact sequence with a modification of the Bloch-Ogus sequence which is still exact. Assume A is a semi-local ring which is smooth and essentially of finite type over a field k ($k = \mathbb{Q}$ in characteristic 0). Consider, as in the proof earlier,

$$(14) \quad \begin{array}{ccccc} 0 \rightarrow K_2(A)\{\ell\} \rightarrow & K_2(F)\{\ell\} \rightarrow & \coprod_{x \in \text{Spec}(A)^1} K_1(\kappa(x))\{\ell\} \\ \downarrow \bar{c}_{2,1}(A) & \downarrow \bar{c}_{2,1}(F) & \downarrow = \\ 0 \rightarrow H^1(A, (2))/\text{Div}^1(A) \rightarrow & H^1(F, (2))/\text{Div}^1(F) \rightarrow & \coprod_{x \in \text{Spec}(A)^1} H^0(\kappa(x), (1)) \end{array}$$

where $H^1(A, (2))$ denotes $H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$, etc. for typographical reasons. If $\text{char}(A)$ is positive, then $H^1(A, \mathbb{Z}_\ell(2))$ is finite and so $H^1(A, \mathbb{Q}_\ell(2))$ which maps onto the maximal divisible subgroup of $H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$ vanishes. Thus the bottom sequence in (14) is just the Bloch-Ogus sequence and so $\bar{c}_{2,1}(A)$ is an isomorphism. If $\text{char}(A) = 0$ and A is essentially of finite type over $\mathbb{Q}$, then $H^0(\kappa(x), \mathbb{Q}_\ell/\mathbb{Z}_\ell(1))$ has no divisible part for any $x \in \text{Spec}(A)^{(1)}$. Thus $\text{Div}^1(A) = \text{Div}^1(F)$, and so the bottom sequence is an exact quotient of the Bloch-Ogus sequence. Thus in either case $\bar{c}_{2,1}(A)$ is an isomorphism.

Now suppose A is an arbitrary semi-local ring of positive characteristic. Since both $K_2(A)$ and $H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$ commute with direct limits of rings we may assume that A is a semi-local ring essentially of finite type over a field. We can then use Theorem 1 since $H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$ is local for the étale topology with respect to closed sets by 1) in Example (1) and $K_2(A)\{\ell\}$ is locally of finite presentation and satisfies the epic hensel pair condition. This latter condition follows since $K_3(A; \mathbb{Z}/\ell^n)$ is local for closed sets by 2) in Example (1) and so ${}_{\ell^n}K_2(A)$, being a quotient, satisfies the epic hensel pair condition. Note that the Corollary to Theorem 1 shows that $K_2(A)\{\ell\}$ is then local for the étale topology with respect to closed sets.

The characteristic 0 case is essentially the same but is complicated by the fact that $H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))/\text{Div}^1(A)$ does not behave nicely with

respect to limits of rings. However Jannsen has shown that

$$H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))/\text{Div}^1(A) \cong H_{\text{cont}}^2(A, \mathbb{Z}_\ell(2)) \{\ell\}.$$

In addition he shows that the sequence,

$$0 \rightarrow \varprojlim^1 H^2(B, \mathbb{Z}/\ell^n(2)) \rightarrow H_{\text{cont}}^2(B, \mathbb{Z}_\ell(2)) \rightarrow \varprojlim H^2(B, \mathbb{Z}/\ell^n(2)) \rightarrow 0,$$

is exact where $\varprojlim^1$ refers to the first left derived functor of the system $\{H^2(B, \mathbb{Z}/\ell^n(2))\}$. Consequently

$$H_{\text{cont}}^2(B^h, \mathbb{Z}_\ell(2)) \cong H_{\text{cont}}^2(A, \mathbb{Z}_\ell(2))$$

if B^h is the henselization with respect to I of a presentation $A \cong B/I$ as a quotient of a semi-local ring B smooth over k . This and the corresponding isomorphism

$$H^1(B^h, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2)) \cong H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2))$$

then allow us to conclude that

$$H_{\text{cont}}^2(B^h, \mathbb{Z}_\ell(2)) \{\ell\} \cong H_{\text{cont}}^2(A, \mathbb{Z}_\ell(2)) \{\ell\}.$$

Now we may apply Theorem to show $\bar{c}_{2,1} : K_2(A)\{\ell\} \rightarrow H_{\text{cont}}^2(A, \mathbb{Z}_\ell(2)) \{\ell\}$ is an isomorphism.

Corollary 4 is obtained by noting that Suslin [19, Corollary 2.8] also proved that $H^1(F, \mathbb{Z}_\ell(2))$ is finite if F is a field of positive characteristic. But if A is a unibranch local ring, e.g. a normal local ring, then $H^1(A, \mathbb{Z}_\ell(2)) \hookrightarrow H^1(F, \mathbb{Z}_\ell(2))$ is also finite. Consequently $H^1(A, \mathbb{Q}_\ell(2)) = 0$, and so we have an isomorphism $H^1(A, \mathbb{Q}_\ell/\mathbb{Z}_\ell(2)) \cong H^2(A, \mathbb{Z}_\ell(2))\{\ell\}$.

APPENDIX A.

In this appendix ζ_n will always stand for a primitive n^{th} root of unity. We wish to prove the following theorem. (Note that this is part of [11, Theorem 1] and is proved without assuming the generalized Kato conjecture.)

Theorem 6. *Let A be a connected, regular semi-local ring with quotient field K , and n an odd positive integer relatively prime to the residue characteristics of A . Then*

$$H^2(A, \mathbb{Z}/n(2)) \xrightarrow{j} H^2(K, \mathbb{Z}/n(2))$$

is injective.

We begin with a series of lemmas to calculate the cohomology of the extension obtained by adjoining an n^{th} root of 1. Fix the following situation in order to describe $\mathbb{Z}/n(k)$ and the action of the cyclotomic character on this module. Fix an integer n such that $n = dm$ and

$m = de$ for some positive integers d, e , and let A be a connected ring with $\zeta_m \in A$ and d a unit in A . Assume $A[\zeta_n]$ is connected. Then

$$X^d - \zeta_m = \prod_{i=1}^d \left(X - \zeta_n (\zeta_m^e)^i \right)$$

in $A[\zeta_n]$, and $A[\zeta_n]$ is a Galois extension of A with cyclic Galois group $G_d := \mathbb{Z}/d = \langle \sigma \rangle \subset (\mathbb{Z}/n)^\times$.

Construct a dictionary for the action of G_d on $\mathbb{Z}/n(k)$ by σ sends $\zeta_n \mapsto \zeta_n (\zeta_m^e)$. Since $n \mid m^2$, we have

G_d on $\mathbb{Z}/n(k)$	$\mathbb{Z}/d$ on $\mathbb{Z}/n$
$(\zeta_n, \zeta_m = \zeta_n^d, \zeta_d = \zeta_m^e) \in \mathbb{Z}/n(1)^{\times 3}$	$(1, d, m = de) \in \mathbb{Z}/n^{\times 3}$
$\sigma(\zeta_n) = \zeta_n (\zeta_m^e) = \zeta_n^{1+m} \in \mathbb{Z}/n(1)$	$\sigma(1) = 1 + m \in \mathbb{Z}/n$
$\sigma^k(\zeta_n) = (\zeta_n^{1+m})^k = \zeta_n^{1+km} \in \mathbb{Z}/n(k)$	$\sigma^k(1) = (1 + m)^k = 1 + km \in \mathbb{Z}/n$

This dictionary makes the computation of the cohomology of G_d on $\mathbb{Z}/n(k)$ straightforward.

Lemma 2. $H^0(G_d, \mathbb{Z}/n(k)) = \mathbb{Z}/(m(d, k))$, and

$$H^{2r}(G_d, \mathbb{Z}/n(k)) = \begin{cases} \mathbb{Z}/(d, k) & \text{if } d \text{ is odd or } k \text{ is even or } 4 \mid m \\ \mathbb{Z}/2(d, k) & \text{if } d \text{ is even and } k \text{ is odd and } 4 \nmid m \end{cases}$$

and

$$H^{2r+1}(G_d, \mathbb{Z}/n(k)) = \begin{cases} \mathbb{Z}/(d, k) & \text{if } d \text{ is odd or } k \text{ is even or } 4 \mid m \\ \mathbb{Z}/2(d, k) & \text{if } d \text{ is even and } k \text{ is odd and } 4 \nmid m \end{cases}$$

Proof. The standard resolution used to calculate the cohomology of a cyclic group G_d on $\mathbb{Z}/n(k)$ is used to carry out this calculation. Our dictionary allows us to calculate $N_k = \sum_{i=0}^{d-1} \sigma^{ik} : \mathbb{Z}/n \rightarrow \mathbb{Z}/n$ and $T_k = 1 - \sigma^k : \mathbb{Z}/n \rightarrow \mathbb{Z}/n$ as

$$\begin{aligned} T_k(j) &= j - j(1 + km) = -kmj \\ N_k(j) &= j \left(\sum_{i=0}^{d-1} (1 + ikm) \right) \\ &= j \left(d + km \frac{d(d-1)}{2} \right) \\ &= \begin{cases} jd & \text{if } d \text{ is odd or } k \text{ is even} \\ j \left(d - m \frac{d}{2} \right) & \text{if } d \text{ is even and } k \text{ is odd} \end{cases} \end{aligned}$$

In the case when d is even and k is odd, we note that

$$\left(m, 1 - \frac{m}{2} \right) = \begin{cases} 2 & \text{if } 4 \nmid m \\ 1 & \text{otherwise} \end{cases}.$$

Then the sequence that calculates the cohomology groups is

$$\cdots \mathbb{Z}/n \xrightarrow{N_k} \mathbb{Z}/n \xrightarrow{T_k} \mathbb{Z}/n \xrightarrow{N_k} \mathbb{Z}/n \xrightarrow{T_k} \mathbb{Z}/n \cdots,$$

and we find

$$\begin{aligned} \text{Ker } T_k &= \frac{d}{(d, k)} \mathbb{Z}/n \cong \mathbb{Z}/(m(d, k)) \\ \text{Im } T_k &= km\mathbb{Z}/n = (d, k) \cdot m\mathbb{Z}/n \\ \text{Ker } N_k &= \begin{cases} m\mathbb{Z}/n & \text{if } d \text{ is odd or } k \text{ is even or } 4 \mid m \\ \frac{m}{2}\mathbb{Z}/n & \text{if } d \text{ is even, } k \text{ is odd, and } 4 \nmid m \end{cases} \\ \text{Im } N_k &= \begin{cases} d\mathbb{Z}/n & \text{if } d \text{ is odd or } k \text{ is even or } 4 \mid m \\ 2d\mathbb{Z}/n & \text{if } d \text{ is even, } k \text{ is odd, and } 4 \nmid m \end{cases} \end{aligned}$$

The rest is straightforward. $\square$

Corollary 6. *Let A be a regular, connected semi-local domain, and let n be an odd positive integer relatively prime to the residue characteristics of A . Suppose that m is the largest divisor of n such that $\zeta_m \in A$. Let $A_n = A[\zeta_n]$, and let $G = \text{Gal}(A_n/A)$. Then $H^p(G, \mathbb{Z}/n(2)) = \begin{cases} \mathbb{Z}/m(2) & \text{if } p = 0 \\ 0 & \text{if } p > 0 \end{cases}$*

Proof. If $m = n$, we are done. Suppose that $n = \ell n'$ where ℓ is the smallest prime divisor of $\frac{n}{m}$. Then there is a cyclic subgroup $H \triangleleft G$ such that $A_{n'} := A^H = A[\zeta_{n'}]$, and $A_{n'}$ is a Galois extension of A with abelian Galois group G/H .

If $\zeta_\ell \in A_{n'}$, then A_n and $A_{n'}$ satisfy the hypothesis of the lemma. Hence

$$H^p(H, \mathbb{Z}/n(2)) = \begin{cases} \mathbb{Z}/n'(2) & \text{if } p = 0 \\ 0 & \text{if } p > 0 \end{cases}.$$

If $\zeta_\ell \notin A_{n'}$, then $[A_n : A_{n'}] = \ell - 1$ is relatively prime to n' . Hence

$$H^p(H, \mathbb{Z}/n(2)) = \begin{cases} \mathbb{Z}/n'(2) & \text{if } p = 0 \\ 0 & \text{if } p > 0 \end{cases}.$$

Now by using the Hochschild-Serre spectral sequence,

$$E_2^{p,q} = H^p(G/H, H^q(H, \mathbb{Z}/n(2))) \Rightarrow H^{p+q}(G, \mathbb{Z}/n(2)),$$

and the above calculations, we conclude that $H^p(G/H, H^0(H, \mathbb{Z}/n(2))) \xrightarrow{\cong} H^p(G, \mathbb{Z}/n(2))$. Since $H^0(H, \mathbb{Z}/n(2)) = \mathbb{Z}/n'(2)$, we may apply an induction hypothesis and thus prove the corollary. $\square$

Lemma 3. *Let A be a regular, connected, semi-local domain with quotient field K , n a positive integer relatively prime to the residue characteristics of A . Let B be a Galois extension of A with quotient field L , and let $G = \text{Gal}(B/A)$. Suppose $\zeta_n \in B$. Then*

$$H^1(G, H^1(B, \mathbb{Z}/n(2))) \rightarrow H^1(G, H^1(L, \mathbb{Z}/n(2)))$$

is injective.

Proof. Since A is a UFD, we have an exact sequence

$$0 \rightarrow A^* \rightarrow K^* \xrightarrow{\Pi^{ord_x}} \coprod_{x \in A^{(1)}} i_{x*} \mathbb{Z} \rightarrow 0$$

where $A^{(1)} = \{x \in \text{Spec}(A) \mid ht(x) = 1\}$ and $ord_x : K^* \rightarrow i_{x*} \mathbb{Z}$ calculates the order of $f \in K^*$ at the discrete valuation associated to x , and $i_x : \text{Spec}(k(x)) \rightarrow \text{Spec}(A)$ is the inclusion of the residue field of A_x . If we map this sequence to itself by multiplication by n , the cokernels form the short exact sequence

$$0 \rightarrow A^*/A^{*n} \rightarrow K^*/K^{*n} \xrightarrow{\Pi^{ord_x}} \coprod_{x \in A^{(1)}} i_{x*} \mathbb{Z}/n \rightarrow 0.$$

There is a sequence of G -modules obtained by twisting the sequence for B with the cyclotomic character

$$0 \rightarrow B^*/B^{*n}(1) \rightarrow L^*/L^{*n}(1) \rightarrow \coprod_{y \in B^{(1)}} i_{y*} \mathbb{Z}/n(1) \rightarrow 0,$$

relating $B^*/B^{*n}(1) = H^1(B, \mathbb{Z}/n(2))$ to $L^*/L^{*n}(1) = H^1(L, \mathbb{Z}/n(2))$ since $\zeta_n \in L^*$. Thus we need only show that

$$H^0(G, L^*/L^{*n}(1)) \rightarrow H^0(G, \coprod_{y \in B^{(1)}} i_{y*} \mathbb{Z}/n(1))$$

is onto to complete the proof of the lemma. But

$$H^0(G, \coprod_{y \in B^{(1)}} i_{y*} \mathbb{Z}/n(1)) \cong \coprod_{x \in A^{(1)}} H^0(D_{y|x}, i_{y*} \mathbb{Z}/n(1)) \cong \coprod_{x \in A^{(1)}} i_{x*} \mathbb{Z}/m_x(1)$$

where $D_{y|x}$ is the decomposition group of a choice of $y \in B^{(1)}$ lying over $x \in A^{(1)}$ and $\mathbb{Z}/m_x(1)$ is the group of roots of unity of order n in $k(y)$ fixed by $D_{y|x}$ (expressed in the above notation).

Any element $\alpha \in \coprod_{x \in A^{(1)}} i_{x*} \mathbb{Z}/m_x(1)$ can be written as

$$\alpha = \sum_{x \in A^{(1)}} \delta_x \cdot \zeta_n^{r_x}$$

where the sum is finite and $\delta_x = 0$ for $x' \in A^{(1)}$ if $x' \neq x$ and $\delta_x = 1 \in i_{x*} \mathbb{Z}/n$ at $x \in A^{(1)}$. Then there is $f_x \in K^*/K^{*n}$ with $ord_x(f_x) = \delta_x$

and $\text{ord}_{x'}(f_x) = 0$ if $x' \neq x$. Hence $\sum_{x \in A^{(1)}} \left(\sum_{\sigma \in G/D_{y|x}} f_x \sigma(\zeta_n^{r_x}) \right)$ has image α . $\square$

Remark 1. *The proof amounts to first reducing to the case of a dvr A by using the UFD sequence and then solving the problem at the completion where $D_{y|x}$ becomes the Galois group and finally spreading that solution around using the transitivity of the action of G on $\{y \in B^{(1)}/y \text{ lies over a fixed } x \in A^{(1)}\}$.*

Proof. If $\zeta_n \in A$, the theorem reduces to the well known result $Br(A) \subset Br(K)$ since $H^2(A, \mathbb{Z}/n(k)) \cong H^2(A, \mathbb{Z}/n(1)) \cong {}_n Br(A)$ and similarly for K . We reduce the theorem to this case by analyzing the cohomology of a cyclic Galois covering $A[\zeta_n]/A$ with Galois group G . There is a spectral sequence for this covering

$$E_2^{p,q} = H^p(G, H^q(A[\zeta_n], \mathbb{Z}/n(2))) \Rightarrow H^n(A, \mathbb{Z}/n(2)).$$

Suppose $x \in \text{Ker}[H^2(A, \mathbb{Z}/n(2)) \rightarrow H^2(K, \mathbb{Z}/n(2))]$. Then $x = 0$ in $H^2(A[\zeta_n], \mathbb{Z}/n(2))^G$ by the above remarks. But Lemma 3 shows that $x = 0$ also in $E_\infty^{1,1} \subseteq H^1(G, H^1(A[\zeta_n], \mathbb{Z}/n(2)))$. Finally Corollary 6 shows that $x \in E_\infty^{2,0} \subseteq H^2(G, \mathbb{Z}/m(2))$ where $\zeta_m \in A$. But $H^2(G, \mathbb{Z}/m(2))$ is the same for the Galois covering $A[\zeta_n]/A$ and $K[\zeta_n]/K$. Hence $x = 0$ and the theorem follows. $\square$

REFERENCES

- [1] S. Bloch and A. Ogus “Gersten’s conjecture and the homology of schemes”, Ann Sci Ecole Norm Sup., 4th sér. 7(1974), 181-202.
- [2] J-L Colliot-Thélène and M. Ojanguren “Espaces Principaux homogènes localement triviaux”, Publications Mathématiques de l’IHES, 75(1992), 97-122.
- [3] J.-L. Colliot-Thélène, R.T. Hoobler, and B. Kahn, “The Bloch—Ogus—Gabber Theorem”, Algebraic K-Theory, Fields Inst. Commun., Vol. 16, Amer. Math. Soc., Providence, RI, 1997, 31–94.
- [4] O. Gabber, “K-theory of henselian local rings and henselian pairs”, a letter from Ofer Gabber to M. Karoubi, 83, AMS Contemp Math, 1989.
- [5] O. Gabber, “Affine analog of the proper base change theorem”, Israel J. of Math., 87(1994), 325-335.
- [6] H. Gillet, “Gersten’s conjecture for the K-theory with torsion coefficients of a discrete valuation ring”, J Algebra, 103(1986), 377-380.
- [7] H. Gillet, and M. Levine, “The relative form of Gersten’s conjecture over a discrete valuation ring: The smooth case”, J Pure Appl Algebra, 46(1987), 59-71.
- [8] H. Gillet, personal communication, July 1995.
- [9] D. Grayson, “Universal exactness in algebraic K-theory”, J Pure Appl Algebra, 36(1985), 139-141.
- [10] U. Jannsen, “Continuous étale cohomology”, Math Ann, 280(1988), 207-245.

- [11] B. Kahn, “Deux théorèmes de comparaison en cohomologie étale; applications”, *Duke Math J*, 69(1993), 137-165.
- [12] J. Milne, Étale Cohomology, Princeton University Press, 1980, Princeton, NJ.
- [13] C. Pedrini, and C. Weibel, “Invariants of real curves”, *Rend Sem Mat Univers Politecn Torino*, 49(1991), 139-173.
- [14] D. Quillen, “Higher Algebraic K-theory I”, Algebraic K-theory I: Higher K-theories, Lecture Notes in Math, 341, Springer-Verlag, Berlin, 1973, 85-147.
- [15] Michel Raynaud, Anneaux Locaux Hensélien, Lecture Notes in Math, 169, Springer-Verlag, Berlin, 1970.
- [16] SGA4:Théorie des Topos et Cohomologie Étale des Schemas, Lecture Notes in Math, 305, Springer-Verlag, Berlin, 1973.
- [17] C. Soulé, “K-théorie des anneaux d’entiers de corps de nombres et cohomologie étale”, *Inventiones math*, 55(1979), 251-295.
- [18] R. Strano, “On the étale cohomology of hensel rings”, *Comm in Alg*, 12, 2195-2211.
- [19] A. Suslin, “Torsion in K_2 of fields”, *K-theory*, 1(1987), 5-29.
- [20] J. Tate, “Relations between K_2 and Galois cohomology”, *Inv Math* 36(1976), 257-274.
- [21] C. Weibel, “Étale Chern classes at the prime 2”, Proceedings of the Lake Louise Conference on Algebraic K-Theory, NATO ASI Series C, 407, Kluwer Academic Publishers, Dordrecht 1993, 249-286.